

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/SE04/001644

International filing date: 10 November 2004 (10.11.2004)

Document type: Certified copy of priority document

Document details: Country/Office: SE
Number: 0303429-5
Filing date: 19 December 2003 (19.12.2003)

Date of receipt at the International Bureau: 26 November 2004 (26.11.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

PGT / SE 2004 / 001644

Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.



(71) Sökande CMC Contrast AB, Lund SE
Applicant (s)

(21) Patentansökningsnummer 0303429-5
Patent application number

(86) Ingivningsdatum 2003-12-19
Date of filing

Stockholm, 2004-11-15

För Patent- och registreringsverket
For the Patent- and Registration Office


Gunilla Larsson

Avgift
Fee

PATENT- OCH
REGISTRERINGSVERKET
SWEDEN

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 666 02 86
08-666 02 86

AWAPATENT AB

Kontor/Handläggare

Stockholm/Kjell Larsson/EAG

CMC Contrast AB

Anställningsnr

Vår referens

SE-21011409

1

MRI CONTRAST MEDIUM COMPOSITION FOR ORAL ADMINISTRATION

FIELD OF THE INVENTION

The present invention relates to magnetic resonance imaging (MRI), more specifically to the oral administration of manganese contrast medium compositions for imaging of the liver and a method for MRI of a liver using the manganese contrast medium composition.

BACKGROUND OF THE INVENTION

Magnetic resonance imaging (MRI) is now well established as a medical diagnostic tool. The ability of the technique to generate high quality images and to differentiate between soft tissues without requiring the patient to be exposed to ionizing radiation has contributed to this success.

Manganese is a well-known paramagnetic contrast medium useful for MRI of soft tissues in the body. When administered intravenously as a contrast medium, manganese may however be teratogenic at clinical dosages. Administered intravenously, manganese is also known to interfere with the normal functioning of the heart by replacement of calcium in the muscular cell of the heart. To avoid this problem Mn bound in chelate complexes has been used; however the chelation prevents or reduces the enhancement of Mn binding with tissues.

EP 0 308 983 A2 thus proposes a contrast medium for parenteral application for imaging of the heart and liver in the form of a non-chelate coordination compound to reduce toxicity. Here Mn is bound in a complex, i.e. a coordination complex, to avoid the toxic effect of free Mn. The proposed compound has the formula $\text{Mn(II)(H}_2\text{O)}_m\text{A}_1\text{A}_2\text{A}_3\text{A}_4^{\text{+a}}(\text{Y,Z})^{\text{-a}}$, wherein A1, A2, A3 and A4 are the same or different amino-substituted carboxylic acid groups having from 2 to 18 carbons; Y and Z are each the same or a different anion of a pharmaceutically acceptable inorganic acid or an organic carboxylic acid having from 2 to 18 carbons; a is the valence of the ions; m, n, o, p and q are each 1 or 0, $(m + n + o + p + q) = 4$; and r is 1, and if Y is a multivalent anion, r is 0 or 1.

EP 0 308 983 A2 discloses organ specifically designed compositions for administration by injection and selected

to carry the coordination compound to the portion of the body to be imaged by the MR device.

5 In order to further reduce the risk of dangerous or even fatal effect on the heart, oral administration of manganese has previously been proposed. A result of the vascularisation of the upper gastrointestinal tract is that orally administered material taken up into the blood from the gut is adsorbed in the liver before passing to
10 the heart. In the case of manganese, absorption by the hepatocytes in the liver prevents cardiotoxic levels of manganese reaching the heart. This hepatocyte uptake of manganese has led to the proposed use of orally administered manganese as a liver imaging contrast
15 medium.

WO 96/05867 discloses a contrast medium composition comprising a physiologically tolerable manganese compound, an uptake promoter and a physiologically
20 tolerable carrier or excipient, having a manganese concentration of at least 0.3 mM or being in a dosage unit form containing at least 300 μ mol manganese, wherein the uptake promoter comprises a physiologically tolerable reducing compound containing an α -hydroxy ketone group, a
25 physiologically tolerable acid containing α - and/or β -hydroxy or amino groups, or a salt thereof, and/or vitamin D. The disclosure clearly indicates that improved uptake of a manganese will be obtained by using an uptake promoter in molar excess over manganese.

30 WO 97/02842 discloses the use of a Mn-contrast medium containing a promoter for oral administration for imaging of the stomach, liver, bile duct and gall bladder. The promoter is in the form of at least one amino acid and/or
35 a vitamin D. However, nor does this document teach any advantage of avoiding substantial formation of coordination compounds between manganese and uptake promoter.

40 SUMMARY OF THE INVENTION

It has now surprisingly been found that by increasing previously taught molar ratios of Mn to amino acid it is possible to improve the liver uptake of manganese, and
45 such improvement is independent of whether vitamin D₃ is present or not.

In this connection it is important to note that even a relatively small increase in the amount of manganese

present in the liver tissue will result in a significant enhancement of the image.

5 In one aspect the present invention provides for the use of a physiologically acceptable manganese (II) compound and an uptake promoter in the form of one or more amino acids for the manufacture of an MRI contrast composition for oral administration and MRI examination of the liver, in a ratio of Mn/promoter higher than that at which
10 coordination compounds between Mn and promoter are formed to a substantial degree.

15 In a second aspect the invention provides for an MRI contrast medium composition for oral administration for examination of the liver, comprising as an active ingredient a physiologically acceptable manganese (II) compound and an uptake promoter comprising one or more amino acids, wherein Mn and the promoter are used in a molar ratio higher than that at which coordination
20 compounds between Mn and promoter are formed to a substantial degree.

25 In a third aspect the invention provides for an MRI contrast medium kit comprising a first container accommodating a physiologically acceptable manganese (II) compound, a second container accommodating an uptake promoter comprising one or more amino acids, and, optionally, instructions for use of the kit, the molar ratio of Mn/promoter being within the range of about 2:3
30 to about 3:1.

35 In a fourth aspect the invention provides for a method for MRI of a mammalian liver using a contrast medium composition as defined above, said method comprising oral administration of an effective amount of a contrast composition as described above to a mammal, including man, in need of such an MRI.

40 DETAILED DESCRIPTION OF THE INVENTION

45 The prior art describes on the one hand the necessity of providing manganese in the form of coordination complexes to avoid toxic effects by injection and, on the other hand, the advantage of using the uptake promoter in a molar excess over the manganese when using oral administration.

The main object of the present invention is to provide contrast medium compositions for oral administration of

manganese for improved imaging of the liver due to increased absorption from the gastro-intestinal tract.

5 The present invention is taking advantage of the novel finding that ratios of manganese/uptake promoter which form coordination complexes are not very effective in regard to adsorption in the liver. Accordingly, another object of the present invention is to provide compositions, wherein manganese and uptake promoter are present in such portions as not to form coordination complexes to any significant extent. Expressed in another way the ratio between manganese and promoter shall be higher than that at which coordination compounds between manganese and promoter are formed to a substantial degree. In this context "substantial" is to be interpreted preferably as major, i.e. leaving more than 50% of the manganese ions non-coordinated and available for absorption.

20 Thus, the present invention is based on the surprising finding that coordination compounds involving manganese and uptake promoter are to be avoided in order that improved adsorption of manganese in the liver will be obtained.

25 In the composition according to the invention, the preferred molar ratio of manganese to uptake promoter is higher than or equal to about 2:3; more preferably higher than or equal to about 1:1, most preferably higher than 30 or equal to about 2:1. The upper limit is preferably at most about 3:1. Thus, a preferable range is from about 2:3 to about 3:1.

35 The preferred dosage of the composition according to the present invention will vary according to a number of factors, such as the age, weight and species of the subject, and the particular uptake promoter used. Conveniently, the dosage of manganese will normally be in the range of from about 25 to about 150 $\mu\text{mol/ kg}$ body weight. Preferably the dosage of manganese will be in a range of from about 50 to about 125 $\mu\text{mol/ kg}$ body weight, and more preferably the dosage of manganese will be in the range of from about 50 to about 100 $\mu\text{mol/ kg}$ body weight.

45 The contrast medium composition according to the invention may comprise a manganese compound together with a mixture of two or more uptake promoters, e.g. a mixture of several amino acids.

Compounds which have been found to be suitable for use as uptake promoters include all physiologically acceptable amino acids. Amino acids which are effective as uptake promoters in the compositions of the invention include all the native amino acids, i.e. alanine, valine, leucine, tryptophan, methionine, isoleucine, proline, phenylalanine, serine, glycine, threonine, cysteine, asparagine, glutamine, tyrosine, aspartic acid, glutamic acid, arginine, lysine and histidine.

A preferred group of amino acids for use as uptake promoters in the compositions of this invention is selected from neutral amino acids including asparagine, and aspartic acid. Particularly preferred amino acids are asparagine, aspartic acid and alanine, especially L-alanine.

The invention is preferably practised while using the amino acid(s) as the only promoter(s), but this does not exclude the use of said amino acid(s) together with any other common promoter(s), e.g. vitamin D₃.

Alanine has been used as an example in the present invention. WO 96/05867 shows a general effect of other amino acids, such as glycine, valine, glutamine, aspartic acid, glutamic acid, lysine, arginine, cysteine and methionine. Decisive for the choice of amino acid or amino acids are the price and the stability as well as the taste, appearance and odour of the amino acid to have a product acceptable to the patients.

Paramagnetic materials, such as manganese ions, may act as either positive or negative MRI contrast agents depending upon a number of factors, including the concentration of the ions at the imaging site and the magnetic field strength used in the imaging procedure. At the concentrations of manganese contemplated for use in the compositions of the invention, the manganese-containing contrast medium will in general function as a positive contrast medium.

The compositions of the invention are particularly suited to use as a dispersion in an aqueous medium. For such a purpose the composition may be administered into the gastrointestinal tract orally or via a gastric tube.

It is possible to formulate the contrast medium immediately or shortly prior to administration by mixing the uptake promoter with the manganese compound.

5 As the contrast medium is to be administered orally a patient can administer it himself. The patient is therefore not obliged to stay in the hospital for several hours before being scanned. He can take the oral medium himself without need for medical assistance.

10 The contrast medium composition of the invention may include components other than the uptake promoter, and the manganese compound, for example conventional pharmaceutical formulation aids, such as wetting agents, 15 buffers, disintegrants, binders, fillers, flavouring agents and liquid carrier media, such as sterile water, water/ethanol etc.

20 The pH of the composition is preferably in the acidic range, e.g. 2 to 7, and while the uptake promoter may itself serve to yield a composition with this pH, buffers or pH adjusting agents may be used.

25 The contrast media may be formulated in any edible/drinkable medium and in conventional pharmaceutically administrable forms, such as tablets, capsules, powders, solutions, dispersions, syrups, etc. Since the contrast medium is to be administered orally, a patient can administer the contrast medium himself prior 30 to scanning.

35 The manganese compound, which for oral administration is preferably dissolved or suspended in water, may for example be in the form of a salt, or may be a mixture of different salts. Particularly preferred are salts in which the manganese is present as Mn(II) rather than Mn(III) since the former has a better adsorption profile and is thus more efficient as an MR contrast medium for the liver.

40 Examples of manganese compounds preferred for use in accordance with the invention include pharmaceutically acceptable salts, e.g. manganese chloride, manganese ascorbate, manganese kojate, manganese salicylate and 45 manganese gluconate, especially the chloride.

The invention will now be further described by examples which illustrate the effectiveness of the uptake of manganese for various concentrations of uptake promoter.

Example 1

5 The study was conducted at Scantox, Hestehavevej 6A, DK-4623 Lille Skensved, Denmark.

10 The objective of the studies was to evaluate the uptake of manganese chloride in rats by measurement of the manganese content in the liver as an comparison example.

15 The experiment was performed in 55 outbred Sprague Dawley rats from Taconic M&B A/S, Denmark. The animals in the study were allocated to 11 groups. The test composition was administered once orally by gavage. Animals in Group 1 (control) were only treated with sterile water. Group 2 received only 100 μmol MnCl_2/kg . The other nine treated groups received concurrent oral doses of 100 μmol MnCl_2/kg (Group 3 to 11) and/or 450, 300 and 150 μmol alanine/kg and/or 20, 40 and 60 IU vitamin D_3/kg in 20 different permutations (table 1).

25 Three hours after dose administration, the animals were killed and the livers were weighed and retained frozen. The samples were subsequently homogenised and analysed for manganese concentrations.

30 The highest group mean amount of manganese was seen in group 5, receiving respectively 150 $\mu\text{mol}/\text{kg}$ alanine and 60 IU vitamin D_3/kg , indicating that a low amount of alanine, i.e. 150 $\mu\text{mol}/\text{kg}$ is increasing the manganese uptake in the liver, compared to groups 3 and 4, receiving 450 and 300 $\mu\text{mol}/\text{kg}$ alanine, respectively and the same amount of vitamin D_3 (see table 1). The results in table 1 show clearly that there is a significant 35 increase in manganese uptake from the gut with decreasing amounts of alanine. For the three subgroups 60, 40 and 20 IU vitamin D_3/kg bodyweight, the best manganese uptake is seen with the lowest alanine amount, i.e. 150 $\mu\text{mol}/\text{kg}$ bodyweight (group 5 compared to 3 and 4, group 8 compared to group 6 and 7, group 11 compared to group 9 and 10). 40 However a vitamin D_3 content of 40 IU/kg bodyweight shows no difference between group 7 and 8, which suggest that in this level of vitamin D_3 there is now significant difference between 300 and 150 $\mu\text{mol}/\text{kg}$ alanine.

45

Table 1A (grouped according to vit. D₃ content)

Group	Treatment			Mol. Ratio Mn:Ala	Liver Mean Mn nmol/g
	MnCl ₂ (μ mol/kg b.w.)	Alanine (μ mol/kg b.w.)	Vitamin D ₃ (IU/kg b.w.)		
1	0	0	0	-	41.4
2	100	0	0	-	44.4
3	100	450	60	1:4.5	61.1
4	100	300	60	1:3	61.8
5	100	150	60	1:1.5	76.1
6	100	450	40	1:4.5	66.4
7	100	300	40	1:3	73.8
8	100	150	40	1:1.5	73.7
9	100	450	20	1:4.5	68.4
10	100	300	20	1:3	69.8
11	100	150	20	1:1.5	72.6

5 Taken together, table 1 shows that when the ratio of Mn:ala is 1:4.5, the Mn liver uptake is enhanced by addition of vit D₃ (group 3, 6 and 9). However when the ratio of Mn:ala is 1:1.5 addition of vit.D₃ in a high amount (i.e. 60 or 40 IU/kg b.w.) is not needed in order to obtain fairly good images (group 5, 8 and 11).

10 Example 2

The study was conducted as described in example 1. However the experiment was performed in 55 outbred Sprague Dawley rats from Taconic M&B A/S, Denmark. The animals in the study were allocated to 11 groups. The test composition was administered orally by gavage. Animals in Group 1 (control) were only treated with sterile water. Group 2 received only 100 μ mol MnCl₂/kg. The other nine treated groups received concurrent oral doses of 100 μ mol MnCl₂/kg (Group 3 to 11), 150, 75 and 37.5 μ mol alanine/kg and 20, 10 and 5 IU vitamin D₃/kg in different permutations (table 2A).

25 Three hours after dose administration, the animals were killed and the livers were weighed and retained frozen. The samples were subsequently homogenised and analysed for manganese concentrations.

30 Manganese concentrations in the liver were higher in groups 3 to 11 compared to Groups 1 and 2. This correlates well with the fact that Groups 1 and 2 received either 0 or 100 μ mol MnCl₂/kg without addition

of uptake promoters. The highest group mean amount of manganese was seen in group 6 to 11, receiving respectively 75 and 37.5 $\mu\text{mol/kg}$ alanine, indicating that a high amount of alanine, i.e. 150 $\mu\text{mol/kg}$ is not increasing the manganese uptake in the liver (group 3 to 5) compared to the alanine amounts provided to group 6 to 11 (see table 2A (or 2B, NB: different grouping) and figure 1). The results clearly show that there is a significant increase in manganese uptake from the gut with the use of promoters wherein the molecular ratio between Mn and amino acid, for example alanine is higher than 1:1.

Table 2A (Grouped according to alanine content)

Group	Treatment			Mol. Ratio Mn:Ala	Liver Mean Mn nmol/g
	MnCl ₂ ($\mu\text{mol/kg}$ b.w.)	Alanine ($\mu\text{mol/kg}$ b.w.)	Vitamin D ₃ (IU/kg b.w.)		
1	0	0	0	-	34.6
2	100	0	0	-	63.4
3	100	150	20	1:1.5	69.9
4	100	150	10	1:1.5	72.7
5	100	150	5	1:1.5	68.4
6	100	75	20	1:0.75	69.2
7	100	75	10	1:0.75	73.8
8	100	75	5	1:0.75	72.0
9	100	37.5	20	1:0.375	72.6
10	100	37.5	10	1:0.375	79.8
11	100	37.5	5	1:0.375	73.1

Table 2B (grouped according to Vit. D₃ content)

Group	Treatment			Mol. Ratio Mn:Ala	Liver Mean Mn nmol/g
	MnCl ₂ (μ mol/kg b.w.)	Alanine (μ mol/kg b.w.)	Vitamin D ₃ (IU/kg b.w.)		
1	0	0	0	-	34.6
2	100	0	0	-	63.4
3	100	150	20	1:1.5	69.9
6	100	75	20	1:0.75	69.2
9	100	37.5	20	1:0.375	72.6
4	100	150	10	1:1.5	72.7
7	100	75	10	1:0.75	73.8
10	100	37.5	10	1:0.375	79.8
5	100	150	5	1:1.5	68.4
8	100	75	5	1:0.75	72.0
11	100	37.5	5	1:0.375	73.1

Example 3

- 5 The study was conducted as described in Ex.1 and 2. However the experiment was performed in 40 male outbred Sprague Dawley rats from Taconic M&B A/S, Denmark. The animals in the study were allocated to 8 groups. The test article was administered once orally by gavage. Animals
- 10 in Group 1 (control) were only treated with sterile water. Group 2 received only 100 μ mol MnCl₂/kg b.w. The other six treated groups received concurrent oral doses of 50 mmol (Group 3) or 100 mmol MnCl₂/kg (Group 4 to 8), 25, 50, 200 and 200 mmol Alanine/kg and 0 or 10 IU
- 15 vitamin D₃/kg in different permutations (see also table 3).

- Manganese concentrations in the liver were higher in groups 2, 4, 5, 6, 7 and 8 compared to Groups 1 and 3.
- 20 This correlates well with the fact that Groups 1 and 3 received the lowest amount of manganese i.e. 0 or 50 μ mol MnCl₂/kg, respectively. The highest group mean amount of manganese was seen in group 5 and 7 (Table 3 and Fig. 3).
- 25 It is worth noting that group 3 and 4 showed a remarkable difference in manganese uptake, 95.3 and 110.3 respectively, even though that the molecular ratio between alanine and manganese was the same in the 2 groups, i.e. 2:1. This shows that the amount of manganese
- 30 in the solution is important for an improved manganese uptake (Table 3 and Fig. 3). 100 μ mol/kg bodyweight is far better than 50 μ mol/kg bodyweight.

Furthermore, the manganese uptake indicates that alanine also has to be provided in a certain amount, i.e. > 25 $\mu\text{mol/kg}$ b.w. since the manganese uptake in group 5 giving 50 $\mu\text{mol/kg}$ alanine is better than in group 6 only provided 25 $\mu\text{mol/kg}$, i.e 128.8 nmol/g and 115.9 nmol/g respectively (Table 3 and figure 4).

In conclusion, Group 5 and 7 (treated with a combination of 100 μmol MnCl_2/kg and 50 μmol alanine/kg and, respectively 10 IU vitamin D_3) was the groups having the highest amount of manganese uptake in the liver. However, also compositions without vitamin D_3 show good results.

Table 3

Group	Treatment			Mol. Ratio Mn:Ala	Liver Mean Mn nmol/g
	MnCl_2 ($\mu\text{mol/kg}$ b.w.)	Alanine ($\mu\text{mol/kg}$ b.w.)	Vitamin D_3 (IU/kg b.w.)		
1	0	0	0	-	45.0
2	100	0	0	-	114.0
3	50	100	0	1:2	95.3
4	100	200	0	1:2	110.3
5	100	50	0	1:0.5	128.8
6	100	25	0	1:0.25	115.9
*7	100	50	10	1:0.5	133.8
8	100	25	10	1:0.25	117.8

* Group 7: 1 animal excluded since no Mn was taken up

Discussion:

The above specific examples clearly indicate the advantage of avoiding substantial formation of coordination compounds between manganese and uptake promoter in order to obtain improved gastro-intestinal uptake of manganese for transportation to the liver following oral administration of the contrast medium composition. This finding is contrary to what could be expected in a consideration of the state of the art.

Claims

1. The use of a physiologically acceptable manganese (II) compound and an uptake promoter in the form of one or more amino acids for the manufacture of an MRI contrast composition for oral administration and MRI examination of the liver, in a ratio of Mn to promoter higher than that at which coordination compounds between Mn and promoter are formed to a substantial degree.
2. The use according to claim 1, wherein the molar ratio of Mn to promoter is at least about 2:3.
3. The use according to claim 2, wherein said ratio is at most about 3:1.
4. The use according to any one of claims 2 and 3, wherein said ratio is higher than about 2:3.
5. The use according to claim 4, wherein said ratio is at least about 1:1.
6. The use according to claim 5, wherein said ratio is at least about 2:1.
7. The use according to any one of the preceding claims, wherein the dosage of manganese is in the range of from about 25 to about 150 $\mu\text{mol/kg}$ body weight.
8. The use according to claim 7, wherein the dosage of manganese is in the range of from about 50 to about 125 $\mu\text{mol/kg}$ body weight.
9. The use according to claim 8, wherein the dosage of manganese is in the range of from about 50 to about 100 $\mu\text{mol/kg}$ body weight.
10. The use according to any one of the preceding claims, wherein the uptake promoter is selected from the group consisting of alanine, valine, leucine, tryptophan, methionine, isoleucine, proline, phenylalanine, serine, glycine, threonine, cysteine, asparagine, glutamine, tyrosine, aspartic acid, glutamic acid, arginine, lysine and histidine.

11. The use according to claim 10, wherein said promoter is selected from neutral amino acids including asparagine and aspartic acid.

5 12. The use according to claim 12, wherein said promoter is L-alanine.

10 13. An MRI contrast medium composition for oral administration for examination of the liver comprising as an active ingredient a physiologically acceptable manganese (II) compound and an uptake promoter comprising one or more amino acids wherein Mn and the promoter are used in a molar ratio higher than that at which coordination compounds between Mn and promoter are formed to a substantial degree.

15 14. A composition according to claim 13, wherein said ratio is as defined in any one of claims 2 to 6.

20 15. A composition according to claim 13 or 14, wherein the dosage of manganese is as defined in any one of claims 7 to 9.

25 16. A composition according to any one of claims 13-15, wherein said uptake promoter is as defined in any one of claims 10-12.

30 17. An MRI contrast medium kit comprising a first container accomodating a physiologically acceptable manganese (II) compound, and a second container accomodating an uptake promoter comprising one or more amino acids, and optionally, instructions for the use of the kit, the molar ratio of Mn to promoter being within the range of about 2:3 to about 3:1.

35 18. A kit according to claim 17, wherein said molar ratio, the dosage of manganese and/or said uptake promoter is (are) as defined in any one of claims 4-12.

40 19. A method for MRI of a mammalian liver using an MRI contrast medium composition according to any one of claims 13-16, said method comprising oral administration of an effective amount of said contrast medium composition.


45

Abstract

5 The use of a physiologically acceptable manganese (II)
 compound and an uptake promoter in the form of one or
 more amino acids for the manufacture of an MRI contrast
 composition for oral administration and MRI examination
 of the liver, in a ratio of Mn to promoter higher than
 that at which coordination compounds between Mn and
 promoter are formed to a substantial degree;
 10 an MRI contrast medium composition for such use;
 and MRI contrast medium kit; and
 a method for imaging of a mammalian liver using such
 contrast medium composition.

5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100

Figure 1







Figure 3



030349-3

22 23 24
 25 26 27
 28 29 30
 31 32 33
 34 35 36
 37 38 39
 40 41 42
 43 44 45
 46 47 48
 49 50 51
 52 53 54
 55 56 57
 58 59 60
 61 62 63
 64 65 66
 67 68 69
 70 71 72
 73 74 75
 76 77 78
 79 80 81
 82 83 84
 85 86 87
 88 89 90
 91 92 93
 94 95 96
 97 98 99
 100 101 102
 103 104 105
 106 107 108
 109 110 111
 112 113 114
 115 116 117
 118 119 120
 121 122 123
 124 125 126
 127 128 129
 130 131 132
 133 134 135
 136 137 138
 139 140 141
 142 143 144
 145 146 147
 148 149 150
 151 152 153
 154 155 156
 157 158 159
 160 161 162
 163 164 165
 166 167 168
 169 170 171
 172 173 174
 175 176 177
 178 179 180
 181 182 183
 184 185 186
 187 188 189
 190 191 192
 193 194 195
 196 197 198
 199 200 201
 202 203 204
 205 206 207
 208 209 210
 211 212 213
 214 215 216
 217 218 219
 220 221 222
 223 224 225
 226 227 228
 229 230 231
 232 233 234
 235 236 237
 238 239 240
 241 242 243
 244 245 246
 247 248 249
 250 251 252
 253 254 255
 256 257 258
 259 260 261
 262 263 264
 265 266 267
 268 269 270
 271 272 273
 274 275 276
 277 278 279
 280 281 282
 283 284 285
 286 287 288
 289 290 291
 292 293 294
 295 296 297
 298 299 300
 301 302 303
 304 305 306
 307 308 309
 310 311 312
 313 314 315
 316 317 318
 319 320 321
 322 323 324
 325 326 327
 328 329 330
 331 332 333
 334 335 336
 337 338 339
 340 341 342
 343 344 345
 346 347 348
 349 350 351
 352 353 354
 355 356 357
 358 359 360
 361 362 363
 364 365 366
 367 368 369
 370 371 372
 373 374 375
 376 377 378
 379 380 381
 382 383 384
 385 386 387
 388 389 390
 391 392 393
 394 395 396
 397 398 399
 400 401 402
 403 404 405
 406 407 408
 409 410 411
 412 413 414
 415 416 417
 418 419 420
 421 422 423
 424 425 426
 427 428 429
 430 431 432
 433 434 435
 436 437 438
 439 440 441
 442 443 444
 445 446 447
 448 449 450
 451 452 453
 454 455 456
 457 458 459
 460 461 462
 463 464 465
 466 467 468
 469 470 471
 472 473 474
 475 476 477
 478 479 480
 481 482 483
 484 485 486
 487 488 489
 490 491 492
 493 494 495
 496 497 498
 499 500 501
 502 503 504
 505 506 507
 508 509 510
 511 512 513
 514 515 516
 517 518 519
 520 521 522
 523 524 525
 526 527 528
 529 530 531
 532 533 534
 535 536 537
 538 539 540
 541 542 543
 544 545 546
 547 548 549
 550 551 552
 553 554 555
 556 557 558
 559 560 561
 562 563 564
 565 566 567
 568 569 570
 571 572 573
 574 575 576
 577 578 579
 580 581 582
 583 584 585
 586 587 588
 589 590 591
 592 593 594
 595 596 597
 598 599 600
 601 602 603
 604 605 606
 607 608 609
 610 611 612
 613 614 615
 616 617 618
 619 620 621
 622 623 624
 625 626 627
 628 629 630
 631 632 633
 634 635 636
 637 638 639
 640 641 642
 643 644 645
 646 647 648
 649 650 651
 652 653 654
 655 656 657
 658 659 660
 661 662 663
 664 665 666
 667 668 669
 670 671 672
 673 674 675
 676 677 678
 679 680 681
 682 683 684
 685 686 687
 688 689 690
 691 692 693
 694 695 696
 697 698 699
 700 701 702
 703 704 705
 706 707 708
 709 710 711
 712 713 714
 715 716 717
 718 719 720
 721 722 723
 724 725 726
 727 728 729
 730 731 732
 733 734 735
 736 737 738
 739 740 741
 742 743 744
 745 746 747
 748 749 750
 751 752 753
 754 755 756
 757 758 759
 760 761 762
 763 764 765
 766 767 768
 769 770 771
 772 773 774
 775 776 777
 778 779 780
 781 782 783
 784 785 786
 787 788 789
 790 791 792
 793 794 795
 796 797 798
 799 800 801
 802 803 8

